

## Regioselective Alkoxysulfonylation of Aryl Methyl Ureas using Alkoxycarbonylsulfonyl Chloride


GOTOH Hideki\* and TANABE Yoo\*<sup>2</sup>

### Abstract

*N*<sup>1</sup>-Aryl-*N*<sup>2</sup>-methylureas (ArN<sup>1</sup>HCON<sup>2</sup>HMe) were regioselectively alkoxy-sulfonylated using alkoxycarbonylsulfonyl chloride (ClSCO<sub>2</sub>R) under two alternative reaction conditions; the reaction proceeded at *N*<sup>2</sup>-position in the absence of base, whereas the use of NaH promoted the reaction at *N*<sup>1</sup>-position.

### I. Introduction

Chlorocarbonylsulfonyl chloride (**1**) and alkoxycarbonylsulfonyl chloride (**2**), prepared from **1** and ROH, are a set of unique and readily available ambident electrophilic reagents (Scheme 1).<sup>1)</sup> Zumach's group originally exploited reagents **1** and **2**, which have been utilized for the synthesis of heterocyclic compounds containing sulfur, oxygen, and nitrogen, and reviewed their impressive highlights.<sup>2)</sup> Another utility was performed for a protecting agent for sulfanyl group in the peptide chemistry.<sup>3)</sup>




As a part of our longstanding interest in *S,N*-containing heterocycles,<sup>4)</sup> we previ-

\* Research Chemist, Master of Science, Global R&D-Material Science of Kao Corporation

\*<sup>2</sup> Professor of Chemistry, Doctor of Science, School of Science and Technology, Kwansei Gakuin University

ously reported regioselective  $\alpha$ -methoxycarbonylsulfonylation of ketones and aldehydes and application to a versatile method for preparing thiazolones, thiadiazinones, and 3-indolethiols.<sup>5)</sup> As a further extension, we report here a couple of regioselective *N*-alkoxycarbonylsulfonylations of *N*<sup>1</sup>-Aryl-*N*<sup>2</sup>-methylureas (ArN<sup>1</sup>HCON<sup>2</sup>HMe) **3** using **2** to afford *N*<sup>2</sup>-alkoxycarbonylsulfonylureas **4** and *N*<sup>1</sup>-isopropoxycarbonylsulfonylureas **5** (Scheme 1).

Scheme 1



## II. Results and Discussion

First, the alkoxycarbonylsulfonylation of **3** using **2** in the absence of bases was examined. The reaction proceeded at the *N*<sup>2</sup>-position with high regioselectivity to give **4** under heating conditions (toluene, 100°C). This result is probably owing to the higher inherent nucleophilicity of N<sup>2</sup> than that of N<sup>1</sup>. Table 1 lists the successful results of the present method using several reagents **2** and substrates **3**, which have a specific substitution pattern (3-CF<sub>3</sub>, 3,4-Cl<sub>2</sub>, and 2-F) expecting for the phenyl urea-type herbicidal activity.<sup>6)</sup> Indeed, these reactions produced the desired compounds **4a-e** each as a sole regioisomer in moderate to good yields. But, particular correlation was not observed between structures of **2** or **3** and the yields.


Table 1

entry	Xn	R	Product	yield/%	entry	Xn	R	Product	yield/%
1	3-CF <sub>3</sub>	Me	<b>4a</b>	45	4	2-F	Et	<b>4d</b>	65
2	3,4-Cl	Et	<b>4b</b>	42	5		<i>i</i> -Pr	<b>4e</b>	57
3		<i>i</i> -Pr	<b>4c</b>	62					

Second, regioselective (region complementary) *N*<sup>1</sup>-sulfonylation was investigated.

A pertinent patent disclosed that a 2-F analogue, *N*<sup>1</sup>-alkoxycarbonylsulfonyl-*N*<sup>2</sup>-(2'-fluorophenyl)-*N*<sup>2</sup>-methylureas, had potent and selective phenyl urea type herbicidal activity.<sup>7)</sup> Thus, we focus our attention on the reaction using the urea **3** (Xn=F). Treatment of **3** with NaH resulted in a regioselective anion formation on *N*<sup>1</sup>-position due to the higher acidity than that of *N*<sup>2</sup>-position. Higher nucleophilic *N*<sup>1</sup>-anion switched the regioselectivity to produce *N*<sup>1</sup>-alkoxysulfonyl-*N*<sup>1</sup>-(2'-fluorophenyl)-*N*<sup>2</sup>-methylureas **5a** in clear contrast to the reaction without bases as mentioned before. Table 2 lists the successful results. Solvent screening showed that DMF was superior than CH<sub>3</sub>CN and THF. Highest yield (53%) was obtained when 1.5 equiv of the urea and 1.6 equiv of NaH was used.

Table 2



entry	urea/eq.	NaH/eq.	solvent	yield/%
1	1.1	1.2	CH <sub>3</sub> CN	trace
2			THF	11
3			DMF	30
4	1.5	1.6		53
5	2.0	2.1		28
6	0.5	0.5		21 <sup>a</sup>

a) Based on ClSCO<sub>2</sub>*i*-Pr

In conclusion, we performed regioselective *N*<sup>1</sup>- and *N*<sup>2</sup>-alkoxysulfonylations of *N*<sup>1</sup>-Aryl-*N*<sup>2</sup>-methylureas. Further investigation of herbicidal activity and its structure-activity relationship of these synthesized compounds is under progress.


### III. Experimental

Flash column chromatography was performed with silica gel Merck 60 (230–400 mesh ASTM). Melting points were determined on a hot stage microscope apparatus (Yanagimoto) and were uncorrected. NMR spectra were recorded on a JEOL DELTA 300 spectrometer, operating at 300 MHz for <sup>1</sup>H NMR and 75 MHz for <sup>13</sup>C NMR. Chemical shifts (δppm) in CDCl<sub>3</sub> were reported downfield from TMS (=0 ppm) for <sup>1</sup>H NMR. For <sup>13</sup>C NMR, chemical shifts on a scale relative to (77.00 ppm) as an internal reference. IR Spectra were recorded on a JASCO FT / IR-5300 spec-

trophotometer. Mass spectra were measured on a JEOL JMS-T 100 LC spectrometer.

### Typical Procedure of $N^2$ -alkoxysulfenylation; exemplified by


#### $N$ -(3'-trifluoromethylphenyl)- $N'$ -methyl- $N'$ -methoxycarbonylsulfenylurea (**4a**)



$\text{ClSCO}_2\text{Me}$  (70 mg, 0.55 mmol) was slowly added to a stirred solution of  $N$ -(2'-trifluoromethylphenyl)- $N'$ -methylurea (109 mg, 0.50 mmol) in toluene at  $100^\circ\text{C}$  under an Ar atmosphere, followed by being stirred at the same temp. for 2 h. The mixture was cooled and concentrated to give crude oil, which was purified by  $\text{SiO}_2$ -column chromatography (hexane:  $\text{AcOEt}$ =15: 1) to give the desired product (**4a**; 70 mg, 45%).

pale yellow oil;  $^1\text{H}$  NMR (300 MHz;  $\text{CDCl}_3$ ):  $\delta$  3.37 (3 H, s), 4.00 (3 H, s), 7.29–7.35 ( $^1\text{H}$ , m), 7.37–7.46 ( $^1\text{H}$ , m), 7.60–7.67 ( $^1\text{H}$ , m), 7.69–7.74 ( $^1\text{H}$ , m), 7.81–7.93 ( $^1\text{H}$ , brs);  $^{13}\text{C}$  NMR (75 MHz;  $\text{CDCl}_3$ ):  $\delta$  40.82, 55.18, 116.28 [ $d, ^3J(^{13}\text{C}, ^{19}\text{F})=3\text{ Hz}$ ], 120.20 [ $d, ^3J(^{13}\text{C}, ^{19}\text{F})=3\text{ Hz}$ ], 122.64, 124.14 [ $d, ^1J(^{13}\text{C}, ^{19}\text{F})=271\text{ Hz}$ ], 129.37, 131.19 [ $d, ^2J(^{13}\text{C}, ^{19}\text{F})=33\text{ Hz}$ ], 138.59, 155.07, 169.01; IR (neat) 3331, 2959, 1740, 1676, 1601, 1541, 1495, 1446, 1336, 1255, 1132  $\text{cm}^{-1}$ .

### Typical Procedure of $N^1$ -alkoxysulfenylation; exemplified by $N$ -(2'-fluorophenyl)- $N$ -isopropoxycarbonylsulfenyl- $N'$ -methylurea (**5a**)




$N$ -(2'-fluorophenyl)- $N'$ -methylurea (126 mg, 0.75 mmol) was slowly added to a stirred solution of NaH (60% in oil) (32 mg, 0.80 mmol) in DMF at  $0$ – $5^\circ\text{C}$  under an Ar atmosphere.  $\text{ClSCO}_2i\text{-Pr}$  (77 mg, 0.50 mmol) was slowly added to the mixture at the same temp., and then warmed up to  $20$ – $25^\circ\text{C}$  during 7 h. Water was added to the mixture, which was extracted three times with  $\text{Et}_2\text{O}$ . The combined organic phase was washed with water, dried ( $\text{Na}_2\text{SO}_4$ ) and concentrated. The obtained crude oil was purified by  $\text{SiO}_2$ -column chromatography (hexane:  $\text{AcOEt}$ =20: 1) to give the desired product (**5a**; 61 mg, 43%).

pale yellow crystals; mp  $53$ – $55^\circ\text{C}$ ;  $^1\text{H}$  NMR (300 MHz;  $\text{CDCl}_3$ ):  $\delta$  1.37 (6 H, d,  $J=6.5\text{ Hz}$ ), 3.29 (3 H, s), 5.10 ( $^1\text{H}$ , sep,  $J=6.5\text{ Hz}$ ), 6.97–7.16 (3 H, m), 8.17–8.26 ( $^1\text{H}$ , m), 11.12–11.20 ( $^1\text{H}$ , br);  $^{13}\text{C}$  NMR (75 MHz;  $\text{CDCl}_3$ ):  $\delta$  21.78, 21.87, 30.69, 71.61, 114.81 [ $d, ^2J(^{13}\text{C}, ^{19}\text{F})=19\text{ Hz}$ ], 121.51, 123.79 [ $d, ^3J(^{13}\text{C}, ^{19}\text{F})=9\text{ Hz}$ ], 124.37 [ $d, ^4J(^{13}\text{C}, ^{19}\text{F})=4\text{ Hz}$ ], 126.61 [ $d, ^3J(^{13}\text{C}, ^{19}\text{F})=9\text{ Hz}$ ], 152.03, 152.98 [ $d, ^1J(^{13}\text{C}, ^{19}\text{F})=242\text{ Hz}$ ], 156.03; IR (KBr) 3221, 2978, 1723, 1620, 1551, 1458, 1356, 1314, 1252, 1186, 1101  $\text{cm}^{-1}$ ; HRMS (ESI) calcd for  $\text{C}_{12}\text{H}_{15}\text{FN}_2\text{O}_3\text{S}$  ( $\text{M}+\text{Na}^+$ ) 309.0685 found 309.0680.

#### $N$ -(3',4'-dichlorophenyl)- $N'$ -methyl- $N'$ -ethoxycarbonylsulfenylurea (**4b**)


Following the procedure for the preparation of **4a**, the reaction of  $N$ -(3',4'-



dichlorophenyl)-*N'*-methylurea (110 mg, 0.50 mmol) with ClSCO<sub>2</sub>Et (77 mg, 0.55 mmol) gave the desired product (**4b**; 68 mg, 42%).

pale yellow crystals; mp 95–97°C; <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>): δ 1.36 (3 H, t, *J* = 7.2 Hz), 3.35 (3 H, s), 4.40 (2 H, q, *J* = 7.2 Hz), 7.24–7.30 (1H, m), 7.32–7.38 (1H, m), 7.63–7.66 (1H, m), 7.72–7.80 (1H, brs); <sup>13</sup>C NMR (75 MHz; CDCl<sub>3</sub>): δ 14.30, 40.88, 65.32, 118.82, 121.21, 126.87, 130.35, 132.66, 137.61, 154.94, 168.27; IR (KBr) 3328, 2990, 2936, 1748, 1663, 1580, 1503, 1387, 1294, 1233, 1134 cm<sup>-1</sup>.

***N*-(3',4'-dichlorophenyl)-*N'*-methyl-*N'*-isopropoxycarbonylsulfonylurea (**4c**)**




Following the procedure for the preparation of **4a**, the reaction of *N*-(3',4'-dichlorophenyl)-*N'*-methylurea (110 mg, 0.50 mmol) with ClSCO<sub>2</sub>*i*-Pr (85 mg, 0.55 mmol) gave the

desired product (**4c**; 104 mg, 62%).

pale yellow crystals; mp 109–111°C; <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>): δ 1.34 (6 H, d, *J* = 6.2 Hz), 3.34 (3 H, s), 5.21 (1H, sep, *J* = 6.2 Hz), 7.23–7.29 (1H, m), 7.31–7.37 (1H, m), 7.61–7.66 (1H, m), 7.71–7.83 (1H, m); <sup>13</sup>C NMR (75 MHz; CDCl<sub>3</sub>): δ 21.85, 40.86, 74.27, 118.86, 121.21, 126.75, 130.29, 132.59, 137.63, 155.03, 167.63; IR (KBr) 3301, 2982, 1734, 1661, 1582, 1522, 1476, 1381, 1296, 1173, 1132 cm<sup>-1</sup>.

***N*-(2'-fluorophenyl)-*N'*-methyl-*N'*-ethoxycarbonylsulfonylurea (**4d**)**




Following the procedure for the preparation of **4a**, the reaction of *N*-(2'-fluorophenyl)-*N'*-methylurea (84 mg, 0.50 mmol) with ClSCO<sub>2</sub>Et (77 mg, 0.55 mmol) gave the desired product (**4d**; 88

mg, 65%).

pale yellow oil; <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>): δ 1.36 (3 H, t, *J* = 7.2 Hz), 3.37 (3 H, s), 4.40 (2 H, q, *J* = 7.2 Hz), 6.95–7.16 (3 H, m), 7.98–8.06 (1H, br), 8.10–8.19 (1H, m); <sup>13</sup>C NMR (75 MHz; CDCl<sub>3</sub>): δ 14.24, 40.82, 65.11, 114.60 [d, <sup>2</sup>*J*(<sup>13</sup>C, <sup>19</sup>F) = 18 Hz], 120.92, 123.55 [d, <sup>3</sup>*J*(<sup>13</sup>C, <sup>19</sup>F) = 8 Hz], 124.48 [d, <sup>4</sup>*J*(<sup>13</sup>C, <sup>19</sup>F) = 3 Hz], 126.64 [d, <sup>3</sup>*J*(<sup>13</sup>C, <sup>19</sup>F) = 8 Hz], 152.57 [d, <sup>1</sup>*J*(<sup>13</sup>C, <sup>19</sup>F) = 241 Hz], 154.90, 168.40; IR (KBr) 3331, 2984, 1742, 1680, 1618, 1595, 1520, 1454, 1318, 1254, 1150 cm<sup>-1</sup>.

***N*-(2'-fluorophenyl)-*N'*-methyl-*N'*-isopropoxycarbonylsulfonylurea (**4e**)**



Following the procedure for the preparation of **4a**, the reaction of *N*-(2'-fluorophenyl)-*N'*-methylurea (84 mg, 0.50 mmol) with ClSCO<sub>2</sub>*i*-Pr (85 mg, 0.55 mmol) gave the desired product (**4e**; 82

mg, 57%).

pale yellow oil; <sup>1</sup>H NMR (300 MHz; CDCl<sub>3</sub>): δ 1.34 (6 H, d, *J* = 6.2 Hz), 3.36 (3 H,

s), 5.22 ( $^1\text{H}$ , sep,  $J=6.2$  Hz), 6.95–7.16 (3 H, m), 7.98–8.07 ( $^1\text{H}$ , br), 8.09–8.18 ( $^1\text{H}$ , m);  $^{13}\text{C}$  NMR (75 MHz;  $\text{CDCl}_3$ ):  $\delta$  21.80, 40.82, 74.04, 114.59 [d,  $^2J(^{13}\text{C}, ^{19}\text{F})=20$  Hz], 120.96, 123.52 [d,  $^3J(^{13}\text{C}, ^{19}\text{F})=9$  Hz], 124.46 [d,  $^4J(^{13}\text{C}, ^{19}\text{F})=3$  Hz], 126.67 [d,  $^3J(^{13}\text{C}, ^{19}\text{F})=9$  Hz], 152.59 [d,  $^1J(^{13}\text{C}, ^{19}\text{F})=241$  Hz], 155.00, 167.81; IR (KBr) 3333, 2982, 2936, 1734, 1678, 1537, 1456, 1318, 1258, 1167, 1092  $\text{cm}^{-1}$ .

### Acknowledgements

This research was partially supported by Grant-in-Aids for Scientific Research on Basic Areas (B) “18350056”, Priority Areas (A) “17035087” and “18037068”, and Exploratory Research “17655045” from MEXT. We are grateful to Dr Ryo Satoh, Sumitomo Chemical Co. Ltd. for the collaboration of this study.

### References and Notes

- 1) Chlorocarbonylsulphenyl chloride (**1**) and alkoxycarbonylsulphenyl chloride (**2**) were commercially available from Tokyo Kasei Kogyo Co., Ltd. and Aldrich Chem. Co., Ltd., respectively. **1** is also easily prepared from  $\text{Cl}_3\text{CSCl}$  and 95%  $\text{H}_2\text{SO}_4\text{-H}_2\text{O}$  in a large scale: E. Muehlbauer and W. Weiss, German Pat. 1233882 (*Chem. Abstr.*, 66, 95023 j, 1967).
- 2) a) Zumack, V. G.; Kühle, E. *Angew. Chem. Int. Ed.*, 82, 63, 1970. As for applications of **1**: b) Pilgram K.; Skiles, R. D. *J. Org. Chem.*, 38, 1578, 1973. c) Baldwin D.; Van den Broek, P. *J. Chem. Soc., Perkin Trans. 1*, 375, 1975. d) Howe, R. K.; Gruner, T. A. Franz, J. E. *J. Org. Chem.*, 42, 1813, 1977. e) Goerdeler J.; Nandi, K. *Chem. Ber.*, 114, 549, 1981. f) Kabashima, S.; Okawara, T.; Yamasaki, T.; Furukawa, M. *J. Heterocycl. Chem.*, 28, 1957, 1991.
- 3) a) Greene, T. W.; Wuts, P. G. M. *Protective Groups in Organic Synthesis*; 4th ed.; Wiley: New York, p. 673, 2007. b) Hiskey, R. G.; Muthukumaraswamy, N.; Vunnam, R. R. *J. Org. Chem.*, 40, 950, 1975. c) Schroll, A. L.; Barany, G. *J. Org. Chem.*, 54, 244 1989.
- 4) a) Tanabe, Y.; Yamamoto, H.; Murakami, M.; Yanagi, K.; Kubota, Y.; Okumura, H.; Sanemitsu, Y.; Suzukamo, G. *J. Chem. Soc., Perkin Trans. 1*, 935 1995. b) Tanabe, Y.; Okumura, H.; Nagaosa, M.; Murakami, M. *Bull. Chem. Soc. Jpn.*, 68, 1467 1995. c) Tanabe, Y.; Nagaosa, M.; Nishii, Y. *Heterocycles*, 33, 2033 1995.
- 5) Sanemitsu, Y.; Kawamura, S.; Tanabe, Y. *J. Org. Chem.*, 57, 1053 1992.
- 6) Monaco, T. J.; Weller, S. C.; Ashton, F. M. *Weed Science: Principle and Practice*; 4th ed.; Wiley: New York, 2002.  $\text{CF}_3$ -compound; Fluorometron<sup>®</sup>. 3-Cl-4-Cl-compound; Diuron<sup>®</sup>.
- 7) Tanabe, Y.; Sato, R.; Hashimoto, S.; Ohno, N. *Jpn. Kokai Tokkyo Koho 84 51259*, 1984; *Chem. Abstr.*, 101, 90610 1984.